

POISSON BRACKETS FOR ORTHOGONAL POLYNOMIALS ON THE UNIT CIRCLE

IRINA NENCIU

ABSTRACT. The recently discovered connection of orthogonal polynomials on the unit circle (OPUC) to the defocusing Ablowitz-Ladik integrable system involves the definition of a Poisson structure on the space of Verblunsky coefficients. In this paper, we compute the complete set of Poisson brackets for the monic orthogonal and the orthonormal polynomials on the unit circle, as well as for the second kind polynomials and the Wall polynomials, thus answering a question posed by Cantero and Simon, [CanSim].

1. INTRODUCTION AND BACKGROUND

Over the past few years, there has been a flurry of activity concerning the defocusing Ablowitz-Ladik system and its connection to the theory of orthogonal polynomials on the unit circle (OPUC). Among other things, this involves the definition of a Poisson structure on the set of Verblunsky coefficients. Let us elaborate: Given a probability measure $d\mu$ on S^1 , the unit circle in $\mathbb{C}$, we can construct an orthonormal system of polynomials, ϕ_n , by applying the Gram-Schmidt procedure to $\{1, z, \dots\}$. These obey a recurrence relation; however, to simplify the formulae, we will present the relation for the monic orthogonal polynomials Φ_n :

$$(1.1) \quad \Phi_{n+1}(z) = z\Phi_n(z) - \bar{\alpha}_n \Phi_n^*(z),$$

$$(1.2) \quad \Phi_{n+1}^*(z) = \Phi_n^*(z) - \alpha_n z \Phi_n(z).$$

Here, $\alpha_n \in \mathbb{C}$ are recurrence coefficients, which are called Verblunsky coefficients, and Φ_n^* denotes the reversed polynomial, $\Phi_n^*(z) = z^n \overline{\Phi_n(\frac{1}{\bar{z}})}$. Notice that $\alpha_n = -\overline{\Phi_{n+1}(0)}$. When $d\mu$ is supported at exactly N points, $\alpha_n \in \mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$ for $0 \leq n \leq N-2$ while α_{N-1} is a unimodular complex number. If the support of $d\mu$ is infinite, then the recurrence formula (1.1) will produce an infinite sequence $\{\alpha_n\}_{n \geq 0}$ of Verblunsky coefficients, all of whom are inside the unit disc $\mathbb{D}$. In both of these cases, there is a 1-1 correspondence between the measure and the sequence of coefficients. For more details on the theory of polynomials orthogonal on the unit circle, we refer the reader to the excellent treatise of B. Simon, [Sim1] and [Sim2].

In the finite case ($d\mu$ supported at N points), one can introduce a Poisson bracket on the space of Verblunsky coefficients, which, for two smooth functions f and g

The author wishes to thank Maria Cantero, Barry Simon, and Takayuki Tsuchida for their useful comments and suggestions. This research was partly supported by NSF grant DMS-0111298 and was done while the author was a member of the School of Mathematics of the Institute for Advanced Study in Princeton.

2000 *Mathematics Subject Classification.* Primary 42C05, Secondary 53D17.

on $\mathbb{D}^{N-1} \times S^1$, is¹

$$(1.3) \quad \{f, g\} = i \sum_{k=0}^{N-2} \rho_k^2 \left[\frac{\partial f}{\partial \bar{\alpha}_k} \frac{\partial g}{\partial \alpha_k} - \frac{\partial f}{\partial \alpha_k} \frac{\partial g}{\partial \bar{\alpha}_k} \right],$$

where $\rho_k = \sqrt{1 - |\alpha_k|^2}$ and, for a complex variable $\alpha = u + iv$, $u, v \in \mathbb{R}$, the partial derivatives are defined as usual by

$$\frac{\partial}{\partial \alpha} = \frac{1}{2} \left[\frac{\partial}{\partial u} - i \frac{\partial}{\partial v} \right], \quad \frac{\partial}{\partial \bar{\alpha}} = \frac{1}{2} \left[\frac{\partial}{\partial u} + i \frac{\partial}{\partial v} \right].$$

If the sequence of Verblunsky coefficients is infinite, one can think of the Poisson bracket as applying to functions which depend only on a finite subset of the coefficients—all the polynomials considered here satisfy this condition.

The purpose of this short note is to compute the Poisson brackets of the monic orthogonal and orthonormal polynomials, thought of as functions of the Verblunsky coefficients, as well as of other polynomials naturally associated to the problem, e.g. second kind polynomials and Wall polynomials (for definitions, see the second part of the Introduction). The first to compute Poisson brackets in the context presented here were, to the best of our knowledge, Nenciu and Simon, [NenSim]. Their purpose was to understand the complete integrability of the defocusing Ablowitz-Ladik (AL) equation with periodic boundary conditions; in the process, they compute certain fairly complicated combinations of the Poisson brackets of the Wall polynomials—see Proposition 2.8. (For the origin of the Ablowitz-Ladik equation, see [AblLad1], [AblLad2]; for recent results on this integrable system, obtained through its connection to OPUC, see for example [Nen1], [KilNen], [GekNen], [CanSim], or the review paper [Nen3].) In a recent paper, [CanSim], Cantero and Simon found all but one of the Poisson brackets for the monic orthogonal and the second kind polynomials. In this paper, we give all the Poisson brackets for the OPs (see Theorem 1); they were derived from the combinations of [NenSim], to which they are equivalent (see Proposition 2.8), but we prove them directly by induction. It was recently brought to our attention that results related to those reported here were obtained in the early 1980s by P. Kulish, [Kul], using r -matrix methods. Together with [KakMug1], [KakMug2], this is one of the first papers which investigate the Poisson bracket used in our work.

Before proceeding to our results, let us give some background on orthogonal polynomials on the unit circle. For more details, we direct the reader to [Sim1] and [Sim2]. From the recurrence relations (1.1) and (1.2) implies that $\|\Phi_n\|_{L^2(d\mu)} = \prod_{j=0}^{n-1} \rho_j$, where $\rho_j = \sqrt{1 - |\alpha_j|^2}$. In particular, this means that $\phi_n = \Phi_n / \prod_{j=0}^{n-1} \rho_j$ and the recurrence relations for the orthonormal polynomials read

$$(1.4) \quad \rho_n \phi_{n+1}(z) = z \phi_n(z) - \bar{\alpha}_n \phi_n^*(z),$$

$$(1.5) \quad \rho_n \phi_{n+1}^*(z) = \phi_n^*(z) - \alpha_n z \phi_n(z).$$

Writing (1.1) and (1.2) in matrix form leads to

$$(1.6) \quad \begin{bmatrix} \Phi_{n+1}(z) \\ \Phi_{n+1}^*(z) \end{bmatrix} = \begin{bmatrix} z & -\bar{\alpha}_n \\ -\alpha_n z & 1 \end{bmatrix} \cdot \begin{bmatrix} \Phi_n(z) \\ \Phi_n^*(z) \end{bmatrix} = \begin{bmatrix} z B_n^*(z) & -A_n^*(z) \\ -z A_n(z) & B_n(z) \end{bmatrix} \cdot \begin{bmatrix} \Phi_0(z) \\ \Phi_0^*(z) \end{bmatrix}.$$

¹Note that the sum defining this Poisson bracket goes only up to $N - 2$ and hence the last Verblunsky coefficient, $\alpha_{N-1} \in S^1$, is a Casimir.

The A_n and B_n are monic polynomials of degree n , and they are known as Wall polynomials. For the proof of the second identity in (1.6), see for example Theorem 3.2.10 of [Sim1]. Furthermore, if λ is a complex unimodular number, and $\{\Phi_n^\lambda\}_{n \geq 0}$ are the monic orthogonal polynomials associated to the (rotated) Verblunsky coefficients $\{\lambda\alpha_n\}_{n \geq 0}$, then the recurrence relations for these polynomials become

$$(1.7) \quad \begin{aligned} \begin{bmatrix} \Phi_{n+1}^\lambda(z) \\ \bar{\lambda}(\Phi_{n+1}^\lambda)^*(z) \end{bmatrix} &= \begin{bmatrix} zB_n^*(z) & -A_n^*(z) \\ -zA_n(z) & B_n(z) \end{bmatrix} \cdot \begin{bmatrix} \Phi_0^\lambda(z) \\ \bar{\lambda}(\Phi_0^\lambda)^*(z) \end{bmatrix} \\ &= \begin{bmatrix} zB_n^*(z) & -A_n^*(z) \\ -zA_n(z) & B_n(z) \end{bmatrix} \cdot \begin{bmatrix} 1 \\ \bar{\lambda} \end{bmatrix}. \end{aligned}$$

The case $\lambda = -1$ is special, and $\Psi_n \equiv \Phi_n^{\lambda=-1}$ are called second kind polynomials. Putting all the relations above together, one can easily obtain the Pinter-Nevai formulae, relating the monic orthogonal and second kind polynomials to the Wall polynomials:

$$(1.8) \quad \Phi_n(z) = zB_{n-1}^*(z) - A_{n-1}^*(z)$$

$$(1.9) \quad \Psi_n(z) = zB_{n-1}^*(z) + A_{n-1}^*(z).$$

or, equivalently,

$$(1.10) \quad A_n(z) = \frac{1}{2z}(\Psi_{n+1}^*(z) - \Phi_{n+1}^*(z))$$

$$(1.11) \quad B_n(z) = \frac{1}{2}(\Psi_{n+1}^*(z) + \Phi_{n+1}^*(z)).$$

The following relations will also be useful farther in our work:

$$(1.12) \quad \Phi_n^\lambda = \frac{1+\bar{\lambda}}{2} \cdot \Phi_n + \frac{1-\bar{\lambda}}{2} \cdot \Psi_n, \quad \Psi_n^\lambda = \frac{1-\bar{\lambda}}{2} \cdot \Phi_n + \frac{1+\bar{\lambda}}{2} \cdot \Psi_n.$$

Notation. In order to keep our notation straight, we specify it here. For a sequence of Verblunsky coefficients $\{\alpha_n\}_{n \geq 0}$, we denote

$$\begin{aligned} \Phi_n(z) &\equiv \Phi_n(z; \alpha_0, \dots, \alpha_{n-1}), \\ \Psi_n(z) &\equiv \Phi_n(z; -\alpha_0, \dots, -\alpha_{n-1}), \\ \Phi_n^\lambda(z) &\equiv \Phi_n(z; \lambda\alpha_0, \dots, \lambda\alpha_{n-1}) \\ \Psi_n^\lambda(z) &\equiv \Phi_n(z; -\lambda\alpha_0, \dots, -\lambda\alpha_{n-1}). \end{aligned}$$

Throughout this paper, we think of these polynomials, as well as of the Wall polynomials A_n and B_n , as functions of the Verblunsky coefficients, with $z \in \mathbb{C}$ a parameter. A simple calculation using the recurrence relations shows that, for any $n \geq 0$:

$$\|\Phi_n\| = \|\Phi_n^*\| = \|\Psi_n\| = \|\Psi_n^*\| = \prod_{j=0}^{n-1} \rho_j,$$

where all the norms are taken in $L^2(d\mu)$, and for $n = 0$ we use the usual convention that the empty product is identically equal to 1. So the orthonormal and normalized second kind polynomials are given by

$$\phi_n = \left(\prod_{j=0}^{n-1} \rho_j^{-1} \right) \cdot \Phi_n \quad \text{and} \quad \psi_n = \left(\prod_{j=0}^{n-1} \rho_j^{-1} \right) \cdot \Psi_n.$$

2. MANY POISSON BRACKETS

The main statement of this note is the following:

Theorem 1. *Let $n \geq 0$ and $d\mu$ a probability measure on the circle whose support contains at least n points (and possibly infinitely many). The Poisson brackets of the monic orthogonal and second kind polynomials associated to this measure are given, for any $z \neq w \in \mathbb{C}$, by*

$$(2.1) \quad \{\Phi_n(z), \Phi_n(w)\} = \{\Psi_n(z), \Psi_n(w)\} = 0,$$

$$(2.2) \quad \{\Phi_n(z), \Phi_n^*(w)\} = \frac{iw}{z-w} (\Phi_n(z)\Phi_n^*(w) - \Phi_n(w)\Phi_n^*(z)),$$

$$(2.3) \quad \{\Psi_n(z), \Psi_n^*(w)\} = \frac{iw}{z-w} (\Psi_n(z)\Psi_n^*(w) - \Psi_n(w)\Psi_n^*(z)),$$

$$(2.4) \quad \begin{aligned} \{\Phi_n(z), \Psi_n(w)\} &= \frac{iw}{z-w} (\Phi_n(z)\Psi_n(w) - \Phi_n(w)\Psi_n(z)) \\ &\quad - \frac{i}{2} (\Phi_n(z) - \Psi_n(z)) (\Phi_n^*(w) + \Psi_n^*(w)), \end{aligned}$$

$$(2.5) \quad \begin{aligned} \{\Phi_n(z), \Psi_n^*(w)\} &= \frac{iw}{z-w} (\Psi_n(z)\Phi_n^*(w) - \Psi_n(w)\Phi_n^*(z)) \\ &\quad + \frac{i}{2} (\Phi_n(z) - \Psi_n(z)) (\Phi_n^*(w) - \Psi_n^*(w)). \end{aligned}$$

Remark 2.1. Note that, by using the invariance of the Poisson bracket under complex conjugation

$$(2.6) \quad \{\bar{f}, \bar{g}\} = \overline{\{f, g\}}$$

and the formula for reversed polynomials, one can compute the remaining Poisson brackets from the ones in Theorem 1. We obtain

$$(2.7) \quad \{\Phi_n^*(z), \Phi_n^*(w)\} = \{\Psi_n^*(z), \Psi_n^*(w)\} = 0,$$

$$(2.8) \quad \begin{aligned} \{\Phi_n^*(z), \Psi_n(w)\} &= \frac{-iz}{z-w} (\Psi_n^*(z)\Phi_n(w) - \Psi_n^*(w)\Phi_n(z)) \\ &\quad - \frac{i}{2} (\Phi_n^*(z) - \Psi_n^*(z)) (\Phi_n(w) - \Psi_n(w)), \end{aligned}$$

and

$$(2.9) \quad \begin{aligned} \{\Phi_n^*(z), \Psi_n^*(w)\} &= \frac{-iw}{z-w} (\Phi_n^*(z)\Psi_n^*(w) - \Phi_n^*(w)\Psi_n^*(z)) \\ &\quad + \frac{i}{2} (\Phi_n^*(z) + \Psi_n^*(z)) (\Phi_n^*(w) - \Psi_n^*(w)). \end{aligned}$$

Also note that, since the left and right-hand sides of the all the formulae given above, (2.1)–(2.9), are continuous in $z, w \in \mathbb{C}$, we can find the Poisson brackets for $z = w$ in the usual fashion. For example, formula (2.5) becomes, for $z = w$,

$$(2.10) \quad \begin{aligned} \{\Phi_n(z), \Psi_n^*(z)\} &= -iz (\Psi_n'(z)\Phi_n^*(z) - \Psi_n(z)(\Phi_n^*)'(z)) \\ &\quad + \frac{i}{2} (\Phi_n(z) - \Psi_n(z)) (\Phi_n^*(z) - \Psi_n^*(z)), \end{aligned}$$

where $'$ denotes the usual complex differentiation.

Remark 2.2. Most of the Poisson brackets from Theorem 1 and Remark 2.1 have already been computed by Cantero and Simon [CanSim]. The Poisson bracket that they didn't compute is (2.5) (and its reverse, (2.8)). With this, we know all the brackets of orthogonal polynomials on the unit circle.

Proof. For $n = 0$, all the polynomials are identically equal to 1, and the relations hold trivially. Assume all the identities hold for some $n \geq 0$. We will prove (2.5) for $n + 1$. All the other relations follow similarly.

First assume that $|\text{supp}(d\mu)| > n + 1$ (and hence $\alpha_0, \dots, \alpha_n \in \mathbb{D}$). Using the recurrence relations, we immediately get

$$\begin{aligned}
 \{\Phi_{n+1}(z), \Psi_{n+1}^*(w)\} &= \{z\Phi_n(z) - \bar{\alpha}_n\Phi_n^*(z), \Psi_n^*(w) + \alpha_n w\Psi_n(w)\} \\
 &= z\{\Phi_n(z), \Psi_n^*(w)\} + \alpha_n z w\{\Phi_n(z), \Psi_n(w)\} \\
 &\quad - \bar{\alpha}_n\{\Phi_n^*(z), \Psi_n^*(w)\} - |\alpha_n|^2 w\{\Phi_n^*(z), \Psi_n(w)\} \\
 &\quad - \{\bar{\alpha}_n, \alpha_n\} w\Phi_n^*(z)\Psi_n(w)
 \end{aligned}
 \tag{2.11}$$

The last identity follows since the n^{th} polynomials depend only on $\alpha_0, \dots, \alpha_{n-1}$, and hence their Poisson brackets with α_n or $\bar{\alpha}_n$ are identically 0. Using the induction hypothesis, we substitute the formulae for the brackets on the right hand side and we get

$$\{\Phi_{n+1}(z), \Psi_{n+1}^*(w)\} = \frac{-iw}{z-w} \cdot T_1 + \frac{i}{2} \cdot T_2 - i\rho_n^2 w\Phi_n^*(z)\Psi_n(w),$$

where

$$\begin{aligned}
 T_1 &= [\Psi_{n+1}(z)\Phi_{n+1}^*(w) - \Psi_{n+1}(w)\Phi_{n+1}^*(z)] \\
 &\quad + (w - z)(|\alpha_n|^2\Phi_n(w)\Psi_n^*(z) + \Psi_n(w)\Phi_n^*(z))
 \end{aligned}$$

and

$$\begin{aligned}
 T_2 &= z(\Phi_n(z) - \Psi_n(z))(\Phi_n^*(w) - \Psi_n^*(w)) - \bar{\alpha}_n(\Phi_n^*(z) + \Psi_n^*(z))(\Phi_n^*(w) - \Psi_n^*(w)) \\
 &\quad - \alpha_n z w(\Phi_n(z) - \Psi_n(z))(\Phi_n(w) + \Psi_n(w)) + |\alpha_n|^2 w(\Phi_n^*(z) - \Psi_n^*(z))(\Phi_n(w) - \Psi_n(w)) \\
 &= [\Phi_{n+1}(z) - \Psi_{n+1}(z)][\Phi_{n+1}^*(w) - \Psi_{n+1}^*(w)] - 2|\alpha_n|^2 w(\Phi_n^*(z)\Psi_n(w) + \Phi_n(w)\Psi_n^*(z)).
 \end{aligned}$$

If we add together all the extra terms, we get

$$\begin{aligned}
 &iw(|\alpha_n|^2\Phi_n(w)\Psi_n^*(z) + \Psi_n(w)\Phi_n^*(z)) - i|\alpha_n|^2 w(\Phi_n^*(z)\Psi_n(w) + \Phi_n(w)\Psi_n^*(z)) \\
 &\quad - i(1 - |\alpha_n|^2)w\Phi_n^*(z)\Psi_n(w) \equiv 0,
 \end{aligned}$$

which proves (2.5) for $n + 1$ and $|\text{supp}(d\mu)| > n + 1$. If $|\text{supp}(d\mu)| = n + 1$, then $\alpha_n \in S^1$ and it is a Casimir for the corresponding Poisson bracket. In particular $\{\bar{\alpha}_n, \alpha_n\} = 0$ in the last term of (2.11). Keeping this in mind, and using the fact that $|\alpha_n| = 1$, one can still run through the calculation above and get the result. By induction, (2.5) holds for all $n \geq 0$, as long as $|\text{supp}(d\mu)| \geq n$.

The proofs of the other relations (2.1)–(2.4) run along the same lines. $\square$

We use the results of Theorem 1 to find the Poisson brackets of the orthonormal and normalized second kind orthogonal polynomials ϕ_n , ψ_n , and their reverses ϕ_n^* and ψ_n^* .

Lemma 2.3. *Let $n \geq 0$, and define $R_n = \prod_{k=0}^{n-1} \rho_k^{-1}$ if $n \geq 1$, $R_0 = 1$, to be the inverse of the norm in $L^2(d\mu)$ of the degree n polynomials. Then*

$$(2.12) \quad -\{R_n, \Phi_n(z)\} = \{R_n, \Psi_n(z)\} = \frac{i}{4R_n}(\Phi_n(z) - \Psi_n(z)) = \frac{i}{4}(\phi_n(z) - \psi_n(z)),$$

and

$$(2.13) \quad \{R_n, \Phi_n^*(z)\} = -\{R_n, \Psi_n^*(z)\} = \frac{i}{4R_n}(\Phi_n^*(z) - \Psi_n^*(z)) = \frac{i}{4}(\phi_n^*(z) - \psi_n^*(w)).$$

Proof. For $0 \geq j \geq n-1$ one has $\{R_n, \alpha_j\} = \frac{i}{2}R_n\alpha_j$, and hence the evolution of the α_j 's under the flow generated by the Hamiltonian R_n ($n > j$) is given by

$$\alpha_j(t) = \alpha_j e^{i \frac{t}{2R_n}}.$$

If we denote $e^{i \frac{t}{2R_n}} = \lambda(t)$, with $|\lambda(t)| = 1$, we obtain using (1.12) that:

$$\begin{aligned} \{R_n, \Phi_n(z)\} &= \left. \frac{d}{dt} \right|_{t=0} \Phi_n(z; \{\lambda(t)\alpha_j\}_{j=0}^{n-1}) = \left. \frac{d}{dt} \right|_{t=0} \left(\frac{1+\overline{\lambda(t)}}{2} \Phi_n(z) + \frac{1-\overline{\lambda(t)}}{2} \Psi_n(z) \right) \\ &= -\frac{i}{4R_n} (\Phi_n(z) - \Psi_n(z)) \end{aligned}$$

The other three Poisson brackets are computed in a similar fashion, using (1.12) and the analogous statements for the reversed polynomials. $\square$

This lemma allows us to compute the Poisson brackets of the normalized OP's:

Proposition 2.4. *If $n \geq 0$ and $|\text{supp}(d\mu)| \geq n+1$, then one has the following Poisson brackets for $z \neq w \in \mathbb{C}$:*

$$(2.14) \quad \{\phi_n(z), \phi_n(w)\} = -\{\psi_n(z), \psi_n(w)\} = \frac{i}{4}(\phi_n(z)\psi_n(w) - \phi_n(w)\psi_n(z)),$$

$$(2.15) \quad \begin{aligned} \{\phi_n(z), \phi_n^*(w)\} &= \frac{i}{4}[\phi_n(z)(\phi_n^*(w) - \psi_n^*(w)) + (\phi_n(z) - \psi_n(z))\phi_n^*(w)] \\ &\quad + \frac{iw}{z-w}(\phi_n(z)\phi_n^*(w) - \phi_n(w)\phi_n^*(z)), \end{aligned}$$

$$(2.16) \quad \begin{aligned} \{\psi_n(z), \psi_n^*(w)\} &= \frac{i}{4}[\psi_n(z)(\psi_n^*(w) - \phi_n^*(w)) + (\psi_n(z) - \phi_n(z))\psi_n^*(w)] \\ &\quad + \frac{iw}{z-w}(\psi_n(z)\psi_n^*(w) - \psi_n(w)\psi_n^*(z)), \end{aligned}$$

$$(2.17) \quad \begin{aligned} \{\phi_n(z), \psi_n(w)\} &= \frac{-iw}{z-w}(\phi_n(z)\psi_n(w) - \phi_n(w)\psi_n(z)) \\ &\quad - \frac{i}{2}(\phi_n(z) - \psi_n(z))(\phi_n^*(w) + \psi_n^*(w)) \\ &\quad + \frac{i}{4}(\phi_n(z)\phi_n(w) - \psi_n(z)\psi_n(w)), \end{aligned}$$

$$(2.18) \quad \begin{aligned} \{\phi_n(z), \psi_n^*(w)\} &= \frac{i}{4}[(\phi_n(z) - \psi_n(z))\phi_n^*(w) - \psi_n(z)(\phi_n^*(w) - \psi_n^*(w))] \\ &\quad - \frac{iw}{z-w}(\psi_n(z)\phi_n^*(w) - \psi_n(w)\phi_n^*(z)). \end{aligned}$$

Remark 2.5. Same as in the case of the monic polynomials (Theorem 1), one can obtain the Poisson brackets for $z = w$ by taking limits as $z \rightarrow w$ in the formulae for $z \neq w$.

Note that the condition on the size of the support of the measure $d\mu$ is more restrictive in this proposition than in Theorem 1. Indeed, if $|\text{supp}(d\mu)| = n$, then one can define the monic polynomial Φ_n by the usual orthogonalization procedure applied to z^n , with respect to $\Phi_0, \dots, \Phi_{n-1}$. But since $\Phi_0, \dots, \Phi_{n-1}$ form a basis in $L^2(d\mu)$, this leads to a monic polynomial of degree n which is identically zero in $L^2(d\mu)$. In particular, one cannot normalize this polynomial, and hence the only orthonormal polynomials that can be correctly defined are $\phi_0, \dots, \phi_{n-1}$.

Proof. The results of this proposition follow by simple calculations from Lemma 2.3 and Theorem 1, since R_n is the normalization constant for both the regular and second kind polynomials, and their reverses. For example, the first Poisson bracket (2.14) can be computed as follows:

$$\{\phi_n(z), \phi_n(w)\} = \{R_n, \Psi_n(w)\}R_n\Phi_n(z) + \{\Phi_n(z), R_n\}R_n\Psi_n(w)$$

since $\{R_n, R_n\}$ and $\{\Phi_n(z), \Phi_n(w)\}$ are identically zero. Therefore, using (2.12) and canceling terms, we obtain (2.14). The other formulae can be obtained similarly. $\square$

Finally, we have the following:

Proposition 2.6. *The Poisson brackets of the Wall polynomials are given, for $z \neq w \in \mathbb{C}$, by*

$$(2.19) \quad \{A_n(z), A_n(w)\} = \{B_n(z), B_n(w)\} = 0,$$

$$(2.20) \quad \{A_n(z), A_n^*(w)\} = iA_n(z)A_n^*(w) - \frac{i}{z-w}(zB_n^*(z)B_n(w) - wB_n^*(w)B_n(z)),$$

$$(2.21) \quad \{B_n(z), B_n^*(w)\} = \frac{iz}{z-w}(A_n(z)A_n^*(w) - A_n(w)A_n^*(z)),$$

$$(2.22) \quad \{A_n(z), B_n(w)\} = \frac{-iw}{z-w}(A_n(z)B_n(w) - A_n(w)B_n(z)),$$

and

$$(2.23) \quad \{A_n(z), B_n^*(w)\} = \frac{iz}{z-w}(A_n(z)B_n^*(w) - A_n(w)B_n^*(z)).$$

Remark 2.7. All the other brackets for Wall polynomials can be obtained from the ones above by taking the reversed polynomials in z and w , and using (2.6). The Poisson brackets for $z = w$ can be obtained from the ones above by taking limits on both sides of the identities (see also Remark 2.1). The six Poisson brackets (2.19)–(2.23) for Wall polynomials are equivalent to the Poisson brackets (2.1)–(2.5).

Proof. The six Poisson brackets in this proposition are equivalent to the six Poisson brackets for monic and second kind orthogonal polynomials from Theorem 1. This equivalence is merely a reflection of the Pinter-Nevai formulae (1.8)–(1.11), relating orthogonal and Wall polynomials. The rest is just simple algebra. $\square$

As explained in the Introduction, the first, to the best of our knowledge, computations of Poisson brackets for polynomials associated to OPUC were done in the work of Nenciu and Simon [NenSim] (see, also, [Sim2] and [Nen2]). The authors were interested in understanding the Ablowitz-Ladik system in the periodic setting, and hence wanted to find Poisson brackets for the discriminant associated to this problem. In terms of Wall polynomials, they have to show that

$$\{B_n(z) + zB_n^*(z), B_n(w) + wB_n^*(w)\} = 0.$$

The main observation is that, since the α 's are separated in the definition of the Poisson bracket, one can use induction in n to prove statements such as the one above. Using the recurrence relations for Wall polynomials immediately shows that one needs to compute certain anti-symmetric combinations of Poisson brackets involving both Wall polynomials (A and B), and their reverses. What they show is the following: For $n \geq 0$, set

$$\begin{aligned} F_n(z, w) &= -i\{A_n^*(z), B_n(w)\}, & R_n(z, w) &= i\{B_n^*(z), B_n(w)\}, \\ S_n(z, w) &= i\{A_n(z), A_n^*(w)\}, & X_n(z, w) &= i\{B_n^*(z), A_n^*(w)\}. \end{aligned}$$

and, for $q \in \mathbb{Z}$ and $z, w \in \mathbb{C} \setminus \{0\}$, define

$$Q_q(z, w) = zw \cdot \frac{z^{q-1} - w^{q-1}}{z - w}.$$

Condensing the statements of Nenciu and Simon (see Propositions 4.5 and 4.7 of [Nen2]), one has

Proposition 2.8. *The following statements hold for all $n \geq 0$, $q \in \mathbb{Z}$, and $z, w \neq 0$:*

$$\begin{aligned}
(\alpha_n, \beta_n) \quad & \{A_n^*(z), A_n^*(w)\} = \{B_n(z), B_n(w)\} = 0 \\
(\gamma_{n,q}) \quad & z^q F_n(z, w) - w^q F_n(w, z) = Q_q(z, w) [A_n^*(z) B_n(w) - A_n^*(w) B_n(z)] \\
(r_{n,q}) \quad & z^q R_n(z, w) - w^q R_n(w, z) = Q_q(z, w) [A_n(z) A_n^*(w) - A_n(w) A_n^*(z)] \\
& z^q S_n(z, w) - w^q S_n(w, z) = -[z^q A_n(z) A_n^*(w) - w^q A_n(w) A_n^*(z)] \\
(s_{n,q}) \quad & + \frac{z^q - w^q}{z - w} [z B_n^*(z) B_n(w) - w B_n^*(w) B_n(z)] \\
(x_{n,q}) \quad & z^q X_n(z, w) - w^q X_n(w, z) = Q_q(z, w) [B_n^*(z) A_n^*(w) - B_n^*(w) A_n^*(z)]
\end{aligned}$$

Proof. The original proof of this proposition goes by induction in n , where the n^{th} induction statement consists of the identities $(\alpha_n) - (x_{n,q})$ for all $q \in \mathbb{Z}$. This makes for a fairly complicated proof.

In fact, the sequence of identities $(\gamma_{n,q})_{q \in \mathbb{Z}}$ is equivalent to formula (2.23). Indeed, take $(\gamma_{n,q})$ for $q = 0$ and 1 and insert the definition of F_n . A simple calculation shows that $Q_0(z, w) \equiv -1$ and $Q_1(z, w) \equiv 0$, and hence

$$\begin{aligned}
\{A_n^*(z), B_n(w)\} - \{A_n^*(w), B_n(z)\} &= -i(A_n^*(z) B_n(w) - A_n^*(w) B_n(z)) \\
z\{A_n^*(z), B_n(w)\} - w\{A_n^*(w), B_n(z)\} &= 0
\end{aligned}$$

Multiplying the first equation by w and subtracting yields

$$(2.24) \quad \{A_n^*(z), B_n(w)\} = \frac{iw}{z-w} (A_n^*(z) B_n(w) - A_n^*(w) B_n(z)),$$

which is equivalent to (2.23). Conversely, consider the combination on the left-hand side of $(\gamma_{n,q})$ for some $q \in \mathbb{Z}$ and insert (2.24). Since the factor

$$A_n^*(z) B_n(w) - A_n^*(w) B_n(z)$$

is anti-symmetric in z and w , the right-hand side of $(\gamma_{n,q})$ emerges from

$$z^q \frac{w}{z-w} - w^q \frac{z}{w-z} \cdot (-1) = zw \frac{z^{q-1} - w^{q-1}}{z-w} \equiv Q_q(z, w).$$

Equally elementary considerations show that $(r_{n,q})$ through $(x_{n,q})$ are equivalent to the corresponding identities from Proposition 2.6. $\square$

REFERENCES

- [AblLad1] M. J. Ablowitz, J. F. Ladik, Nonlinear differential-difference equations. *J. Math. Phys.* **16** (1975), 598–603.
- [AblLad2] M. J. Ablowitz, J. F. Ladik, Nonlinear differential-difference equations and Fourier analysis. *J. Math. Phys.* **17** (1976), 1011–1018.
- [CanSim] M. J. Cantero, B. Simon, Poisson brackets of orthogonal polynomials, preprint, [arXiv:math.SP/0610989](https://arxiv.org/abs/math.SP/0610989).
- [GekNen] M. Gekhtman, I. Nenciu, Multi-Hamiltonian structures for the finite defocusing Ablowitz-Ladik system, preprint.
- [KakMug1] F. Kako, N. Mugibayashi, Complete integrability of general nonlinear differential-difference equations solvable by the inverse method. I. *Progr. Theoret. Phys.* **60** (1978), no. 4, 975–984.
- [KakMug2] F. Kako, N. Mugibayashi, Complete integrability of general nonlinear differential-difference equations solvable by the inverse method. II. *Progr. Theoret. Phys.* **61** (1979), no. 3, 776–790.
- [KilNen] R. Killip, I. Nenciu, CMV: the unitary analogue of Jacobi matrices, preprint, to appear in *Comm. Pure Appl. Math.*

- [Kul] P. P. Kulish, Quantum difference nonlinear Schrödinger equation, *Lett. Math. Phys.* **5** (1981), no. 3, 191–197.
- [Nen1] I. Nenciu, Lax pairs for the Ablowitz-Ladik system via orthogonal polynomials on the unit circle, *Int. Math. Res. Not.* **11** (2005), 647–686.
- [Nen2] I. Nenciu, Lax Pairs for the Ablowitz-Ladik System via Orthogonal Polynomials on the Unit Circle, Ph.D. Thesis, Caltech, May 2005.
- [Nen3] I. Nenciu, CMV matrices in random matrix theory and integrable systems: a survey, *J. Phys. A: Math. Gen.* **39** (2006), 8811–8822.
- [NenSim] I. Nenciu, B. Simon, unpublished, has appeared in Section 11.11 of [Sim2] and, with complete proofs, in [Nen2].
- [Sim1] B. Simon, *Orthogonal Polynomials on the Unit Circle, Part 1: Classical Theory*, AMS Colloquium Series, American Mathematical Society, Providence, RI, 2005.
- [Sim2] B. Simon, *Orthogonal Polynomials on the Unit Circle, Part 2: Spectral Theory*, AMS Colloquium Series, American Mathematical Society, Providence, RI, 2005.

IRINA NENCIU, COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012
E-mail address: `nenciu@cims.nyu.edu`